

Effective Versions of the Chebotarev Density Theorem

J.C. Lagarias and A.M. Odlyzko



Typeset by the TeXromancers[†], credits go to: Aareyan Manzoor, Andrew Lin, Celeste Yuan, Seewoo Lee, Sadanand Abhyankar

Contents

Contents	1
1 Introduction	2
2 Outline of the main argument	7
3 Artin L-functions and Mellin transform	9
4 Reduction to the case of Hecke L-functions	15
5 Density of zeroes of Hecke L-functions	17
6 The contour integral	23
7 The explicit formula	28
8 Zero-free regions	32
9 Final estimates	35
Bibliography	40

[†]Link to Github for TeX: <https://github.com/AareyanManzoor/Lagarias-Odlyzko>
A dyslexic friendly version is linked in the Github.

1 Introduction

Let K be an algebraic number field (finite extension of the rationals \mathbb{Q}) and L a normal extension of K with Galois group $G = G(L/K)$. Let Δ_L and Δ_K denote the absolute values of the discriminants of L and K , respectively, and let $n_L = [L : \mathbb{Q}]$, $n_K = [K : \mathbb{Q}]$. Throughout this paper \mathfrak{p} will denote a prime ideal of K and P a prime ideal of L . If \mathfrak{p} is a prime ideal of K which is unramified in L , then we use the Artin symbol $\left[\frac{L/K}{\mathfrak{p}} \right]$ to denote the conjugacy class of Frobenius automorphisms corresponding to prime ideals $P \mid \mathfrak{p}$. For each conjugacy class C of G , we define

$$\pi_C(x, L/K) = \left| \left\{ \mathfrak{p} : \mathfrak{p} \text{ unramified in } L, \left[\frac{L/K}{\mathfrak{p}} \right] = C, \mathbb{N}_{K/\mathbb{Q}} \mathfrak{p} \leq x \right\} \right|.$$

The Chebotarev density theorem [Tsc26] asserts that

$$\pi_C(x, L/K) \sim \frac{|C|}{|G|} \text{Li}(x) \quad \text{as } x \rightarrow \infty, \quad (1.1)$$

where $\text{Li}(x)$ is the familiar logarithmic integral

$$\text{Li}(x) = \int_2^x \frac{dt}{\log t} \sim \frac{x}{\log x} \quad \text{as } x \rightarrow \infty$$

The Chebotarev density theorem generalizes many of the classical results on the distribution of primes and prime ideals. For example, if we consider the trivial extension $L = K$ of K (K does not have to be normal over \mathbb{Q}), then there is only one conjugacy class, and (1.1) shows that the number of prime ideals of K with norm $\leq x$ is asymptotic to $\text{Li}(x)$, which is exactly the prime ideal theorem. If we let $K = \mathbb{Q}$ and $L = \mathbb{Q}(e^{2\pi i/q})$, then the conjugacy classes of G correspond to the residue classes modulo q , and (1.1) gives us the prime number theorem for arithmetic progressions.

One of the most important of the many applications of the Chebotarev density theorem deals with the group of an equation. Suppose that $f(x)$ is a monic polynomial whose coefficients are algebraic integers in K and which is irreducible over K . Suppose further that L is the splitting field of $f(x)$ over K . If we regard $G = G(L/K)$ as a permutation group acting on the roots of $f(x)$, then for almost all prime ideals \mathfrak{p} of K the cycle structure of $\left[\frac{L/K}{\mathfrak{p}} \right]$ depends on the factorization of $f(x)$ modulo \mathfrak{p} , and vice versa. Thus if G is known, then the Chebotarev density theorem tells us how often various factorizations occur as \mathfrak{p} runs through all the prime ideals of K . On the other hand, if we do not know G , then factoring $f(x)$ modulo the prime ideals of K will yield the complete cycle structures of G , since by (1.1) for every conjugacy class C there are infinitely many primes \mathfrak{p} with $\left[\frac{L/K}{\mathfrak{p}} \right] = C$. This can be very helpful in the determination of G [Wae70, vol. 1, pp. 189-192], especially since by considering enough primes we can even determine the relative densities of elements of G which have a given cycle structure. (Unfortunately, sometimes this is not enough to determine G completely, since it is possible to construct two nonisomorphic groups which have transitive permutation representations in which the number of elements with a given cycle structure is the same for both groups.) In these situations it is important to be able to compute a bound below which every conjugacy class will occur as the Artin symbol of a prime ideal of K .

The usual proofs of the Chebotarev theorem contains either no error estimates at all, or else estimates which contain constants depending in some undetermined way on the fields K and L . In particular, such estimates do not allow us to specify effectively a value $x_0 = x_0(L/K)$ such that

$$\pi_C(x, L/K) > 0 \quad \text{if } x \geq x_0. \quad (1.2)$$

The purpose of this paper is to prove two versions of the Chebotarev theorem,

each of which has an error term which is an explicit and effectively computable function of x , n_L , Δ_L , and $|C|/|G|$. One version assumes the truth of the Generalized Riemann Hypothesis (GRH) and the other holds unconditionally.

We first state the conditional result.

Theorem 1.1. There exists an effectively computable positive absolute constant c_1 such that if GRH holds for the Dedekind zeta function of L , then for every $x > 2$,

$$\left| \pi_C(x, L/K) - \frac{|C|}{|G|} \operatorname{Li}(x) \right| \leq c_1 \left\{ \frac{|C|}{|G|} x^{\frac{1}{2}} \log(\Delta_L x^{n_L}) + \log \Delta_L \right\}. \quad (1.3)$$

This theorem yields immediately a value of x_0 such that (1.2) holds. (We utilize here the estimate $n_L^{-1} \log \Delta_L > 1 + \varepsilon$ for some $\varepsilon > 0$, valid for $n_L > 1$. It follows from Minkowski's discriminant bound, and it can also be derived from (5.11) (see [Odl77]).

Corollary 1.2. There exists an effectively computable positive absolute constant c_2 such that if the GRH holds for the Dedekind zeta function of $L \neq \mathbb{Q}$, then for every conjugacy class of G there exists an unramified prime ideal \mathfrak{p} in K such that $\left[\frac{L/K}{\mathfrak{p}} \right] = C$ and

$$N_{K/\mathbb{Q}} \mathfrak{p} \leq c_2 (\log \Delta_L)^2 (\log \log \Delta_L)^4. \quad (1.4)$$

(If $L = \mathbb{Q}$, $\mathfrak{p} = (2)$ yields a solution.)

At the end of this paper we will indicate how the above estimate can be improved so as to eliminate the $\log \log \Delta_L$ term.

We next state the unconditional result.

Theorem 1.3. If $n_L > 1$ then $\zeta_L(s)$ has at most one zero in the region defined

by $s = \sigma + it$ with

$$1 - (4 \log \Delta_L)^{-1} \leq \sigma \leq 1, \quad |t| \leq (4 \log \Delta_L)^{-1}. \quad (1.5)$$

(If $n_L = 1$, $L = \mathbb{Q}$ and there is no zero in $|t| \leq 14$, $\sigma > 0$.)

If such a zero exists, it must be real and simple, and we denote it by β_0 .

Further, there exist absolute effectively computable constant c_3 and c_4 such that if

$$x \geq \exp(10n_L(\log \Delta_L)^2), \quad (1.6)$$

then

$$\left| \pi_C(x) - \frac{|C|}{|G|} \text{Li}(x) \right| \leq \frac{|C|}{|G|} \text{Li}(x^{\beta_0}) + c_3 x \exp(-c_4 n_L^{-\frac{1}{2}} (\log x)^{\frac{1}{2}}), \quad (1.7)$$

where the β_0 term is present only when β_0 exists.

Because of the presence of the β_0 factor, Theorem 1.3 does not fully meet our criterion of effectiveness, which is that the error term should depend only on x , n_L , Δ_L , and $|C|/|G|$. However, this defect can be remedied by utilizing any effective bound for β_0 . In most cases the best known such bound is that of Stark [Sta74, p.148] which we quote below.

Theorem 1.4. Let the notation be as in Theorem 1.3, and let $m_L = 4$ if 1 is normal over \mathbb{Q} , $m_L = 16$ if there is a sequence of fields

$$\mathbb{Q} = k_1 \subset k_2 \subset \cdots \subset k_r = L$$

with each field normal over the preceding one, and $m_L = 4n_L!$ otherwise. Then there exists an effectively computable absolute constant c_5 such that

$$\beta_0 < \max[1 - (m_L \log \Delta_L)^{-1}, 1 - (c_5 \Delta_L^{1/n_L})^{-1}]. \quad (1.8)$$

Even if β_0 does not exist, Theorem 1.3 does not give a good unconditional bound for the smallest norm of a prime ideal whose Artin symbol is a given conjugacy class. A reasonable conjecture might be that there should be an effectively computable absolute constant c such that for every normal extension L/K and every conjugacy class C of $G(L/K)$, there should be an unramified \mathfrak{p} with $\left[\frac{L/K}{\mathfrak{p}}\right] = C$ and

$$N_{K/\mathbb{Q}} \mathfrak{p} \leq (\log \Delta_L)^c. \quad (1.9)$$

When L is a cyclomic extension of $K = \mathbb{Q}$, (1.9) is equivalent to Linnik's theorem [Bom74, p.39]. However, if $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{d})$ is a quadratic extension of \mathbb{Q} , the determination of the least prime p with $\left[\frac{L/\mathbb{Q}}{(p)}\right] \neq \{1\}$ corresponds to the problem of determining the least quadratic nonresidue (mod d), and for this problem no unconditional bound better than

$$p \leq c_6 \Delta_L^{c_7} \quad (1.10)$$

is known, where c_6 and c_7 are positive constants. Thus without some major new ideas it would probably be very difficult to prove an unconditional result as good as (1.9). However, by using slightly different techniques (which are designed to detect prime ideals rather than estimate their total number) one can prove the following result [LMO79].

Theorem. There exist effectively computable positive absolute constants b_1 and b_2 such that for every conjugacy class C of G there exists an unramified prime ideal \mathfrak{p} of K such that $\left[\frac{L/K}{\mathfrak{p}}\right] = C$ and

$$N_{K/\mathbb{Q}} \mathfrak{p} \leq b_1 \Delta_L^{b_2}.$$

The approach used in this paper has a long history. The argument given here

may be viewed as a direct descendent of de la Vallée Poussin's proof of the prime number theorem. We follow closely with the pattern of Davenport's treatment [DM13] of the prime number theorem for arithmetic progressions. The main innovation here is the careful treatment of the dependencies of various constants on n_L and Δ_L (cf. [DM13]; [Fog61]; [Gol70]; [Lan71]; [Mor]).

Aside from some slight acquaintance with algebraic and analytic number theory, this paper also assumes knowledge of the basic properties of Hecke and Artin L -functions [Hei67]. The deepest of these results is the abelian reciprocity law, which tells us that an abelian Artin L -series is a Hecke L -series, and so is analytic for $s \neq 1$.

Throughout this paper c_1, c_2, \dots will denote effectively computable positive absolute constants. (In particular, they are independent of K and L .) The Vinogradov notation

$$f \ll g$$

will be used to denote the existence of an effectively computable positive absolute constant A (not necessary the esame in each occurrence) such that

$$|f| \leq Ag,$$

in the range indicated.

2 Outline of the main argument

The main argument is primarily concerned with the derivation of an asymptotic formula with an explicit error term of a weighted prime-power-counting function

$\psi_C(x) = \psi_C(x, L/K)$ associated to $\pi_C(x, L/K)$. It is defined by

$$\psi_C(x, L/K) = \sum_{\substack{N_{K/\mathbb{Q}} \mathfrak{p}^m \leq x \\ \mathfrak{p} \text{ unramified} \\ \left[\frac{L/K}{\mathfrak{p}} \right]^m = C}} \log(N_{K/\mathbb{Q}} \mathfrak{p})$$

The details of this argument are complicated, the main steps are simple in conception:

- (i) $\psi_C(x)$ differs from a truncate inverse Mellin transform

$$I_C(x, T) = \frac{1}{2\pi i} \int_{\sigma_0 - iT}^{\sigma_0 + iT} F_C(s) \frac{x^s}{s} ds,$$

by a remainder term $R_1(x, T)$.

- (ii) $F_C(s)$ can in fact be written as a linear combination of logarithmic derivatives of Hecke (abelian) L-functions. As a consequence, all the singularities of $F_C(s)$, which are simply poles only, occur at zeroes and pole of $\zeta_L(s)$.

- (iii) $I_C(x, T)$ differs from a certain contour integral

$$B_C(x, T) = \frac{1}{2\pi i} \oint_{B_T} F_C(s) \frac{x^s}{s} ds,$$

by a remainder term $R_2(x, T)$. This step is traditionally labelled “shifting the line of integration to the left.”

Certain results on the density of zeros of $\zeta_L(s)$ in the critical strip $0 < \operatorname{Re} s < 1$ are necessary to estimate $R_2(x, T)$.

- (iv) The contour integral $B_C(x, T)$ is evaluated by Cauchy’s residue theorem. The integrand has poles at the zeroes and the pole of $\zeta_L(s)$, and the result is a main term $\frac{|C|}{|G|}x$ coming from the pole of $\zeta_L(s)$ at $s = 1$, together with

a certain sum $S(x, T)$ over the zeroes of $\zeta_L(s)$ within the contour B_T .

The end result of these steps is a truncated “explicit formula” for $\psi_C(x)$ with an unconditional error term, which is stated as Theorem 7.1

- (v) The sum over the zeroes $S(x, T)$ is estimated. It is at this point that unproved hypothesis about the zeroes can be helpful. An unconditional upper bound for $|S(x, T)|$ is obtained using the existence of a zero-free region of $\zeta_L(s)$ near the vertical line $\sigma = 1$. A much better estimate for $|S(x, T)|$ is made assuming the Generalized Riemann hypothesis for $\zeta_L(s)$.
- (vi) The asymptotic formula $\psi_C(x) \sim \frac{|C|}{|G|}x$ with an explicit remainder term is derived by making an appropriate choice of T as a function of x , to minimize the accumulated error terms. (This choice depends on whether the GRH is assumed or not, of course.)
- (vii) The asymptotic formula $\pi_C(x) \sim \frac{|C|}{|G|} \text{Li}(x)$ with an explicit remainder term is derived by partial summation from that for $\psi_C(x)$.

The remaining sections of this paper carry out the details (although we will not follow this outline exactly).

3 Artin L-functions and Mellin transform

In this section we establish the relation between $\psi_C(x)$ and a certain truncated inverse Mellin transform. Throughout this and subsequent sections we will use the abbreviations $\pi_C(x)$, $\psi_C(x)$ and N for $\pi_C(x, L/K)$, $\psi_C(x, L/K)$, and $N_{K/\mathbb{Q}}$, respectively. We will also use ϕ to denote irreducible characters of $G = G(L/K)$.

For each irreducible character ϕ of G we define

$$\phi_K(\mathfrak{p}^m) = \frac{1}{e} \sum_{\alpha \in I} \phi(\tau^m \alpha), \quad (3.1)$$

where I is the inertia group of P , one of the prime ideal factors of \mathfrak{p} , $e = |I|$ and τ is one of the Frobenius automorphisms corresponding to \mathfrak{p} . If $L(s, \phi, L/K)$ is the Artin L-series associated to ϕ , then for $\operatorname{Re}(s) > 1$ we have

$$-\frac{L'}{L}(s, \phi, L/K) = \sum_{\mathfrak{p}} \sum_{m=1}^{\infty} \phi_K(\mathfrak{p}^m) \log(\mathbf{N} \mathfrak{p})(\mathbf{N} \mathfrak{p})^{-ms} \quad (3.2)$$

where the outer sum is over all the prime ideals of K . We should also note that the definitions (3.1) and (3.2) apply equally well to reducible characters.

To single out those \mathfrak{p}^m with $\left[\frac{L/K}{\mathfrak{p}}\right]^m = C$, we will use the characters ϕ . (unfortunately this works only to the extent that some extraneous prime powers \mathfrak{p}^m corresponding to \mathfrak{p} that ramify in L are also included.) Suppose that $g \in C$. We define a function $f_C : G \rightarrow \mathbb{C}$ by

$$f_C = \sum_{\phi} \bar{\phi}(g)\phi. \quad (3.3)$$

Then the orthogonality relations for characters imply that

$$f_C(\tau) = \begin{cases} \frac{|G|}{|C|} & \text{if } \tau \in C, \\ 0 & \text{if } \tau \notin C. \end{cases} \quad (3.4)$$

Hence if

$$F_C(s) = -\frac{|C|}{|G|} \sum_{\phi} \bar{\phi}(g) \frac{L'}{L}(s, \phi, L/K), \quad (3.5)$$

then (3.2) through (3.5) show that for $\operatorname{Re}(s) > 1$ we have the Dirichlet series expansion

$$F_C(s) = \sum_{\mathfrak{p}} \sum_{m=1}^{\infty} \theta(\mathfrak{p}^m) \log(\mathbf{N} \mathfrak{p})(\mathbf{N} \mathfrak{p})^{-ms}, \quad (3.6)$$

where for \mathfrak{p} unramified in L we have

$$\theta(\mathfrak{p}^m) = \begin{cases} 1 & \text{if } \left[\frac{L/K}{\mathfrak{p}} \right]^m = C, \\ 0 & \text{otherwise,} \end{cases}$$

and $|\theta(\mathfrak{p}^m)| \leq 1$ if \mathfrak{p} ramifies in L .

Equation (3.6) shows that except for the ramified prime factors, $\psi_C(x)$ is a partial sum of the coefficients of $F_C(s)$. To obtain $\psi_C(x)$ from $F_C(s)$ we will use the following well-known truncated version of the inverse Mellin transform [Tsc26, p. 54], [DM13, pp. 109-110].

Lemma 3.1. If $y > 0$, $\sigma > 0$, and $T > 0$, then

$$\begin{aligned} \left| \frac{1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{y^s}{s} ds - 1 \right| &\leq y^\sigma \min(1, T^{-1} |\log y|^{-1}) && \text{if } y > 1 \\ \left| \frac{1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{y^s}{s} ds - \frac{1}{2} \right| &\leq \sigma T^{-1} && \text{if } y = 1 \end{aligned}$$

and

$$\left| \frac{1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{y^s}{s} ds \right| \leq y^\sigma \min(1, T^{-1} |\log y|^{-1}) \quad \text{if } 0 < y < 1.$$

Let $\sigma_0 > 1$, $x \geq 2$, and define

$$I_C(x, T) = \frac{1}{2\pi i} \int_{\sigma_0-iT}^{\sigma_0+iT} F_C(s) \frac{x^s}{s} ds. \quad (3.7)$$

Since the Dirichlet series in (3.6) is absolutely convergent for $\operatorname{Re}(s) > 1$, we can integrate term by term (with the help of Lemma 3.1) to obtain

$$\left| I_C(x, T) - \sum_{\substack{\mathfrak{p}, m \\ N \mathfrak{p}^m \leq x}} \theta(\mathfrak{p}^m) \log N \mathfrak{p} \right| \leq \sum_{\substack{\mathfrak{p}, m \\ N \mathfrak{p}^m = x}} \{ \log N \mathfrak{p} + \sigma_0 T^{-1} \} + R_0(x, T), \quad (3.8)$$

where

$$R_0(x, T) = \sum_{\substack{\mathfrak{p}, m \\ N\mathfrak{p}^m \neq x}} \left(\frac{x}{N\mathfrak{p}^m} \right)^{\sigma_0} \min \left(1, T^{-1} \left| \log \frac{x}{N\mathfrak{p}^m} \right|^{-1} \right) \log N\mathfrak{p} \quad (3.9)$$

and where the sum on the right side of (3.8) is present only when there are \mathfrak{p} and m with $N\mathfrak{p}^m = x$. Now the sum of the left side of (3.8) equals $\psi_C(x)$, except for the ramified prime terms. However, $N\mathfrak{p} \geq 2$ for each prime ideal \mathfrak{p} , all the ramified prime ideals \mathfrak{p} divide the discriminant of L over K , and so

$$\begin{aligned} \left| \sum_{\substack{\mathfrak{p}, m \\ N\mathfrak{p}^m \leq x}} \theta(\mathfrak{p}^m) \log N\mathfrak{p} - \psi_C(x) \right| &\leq \sum_{\substack{\mathfrak{p}, m \\ \mathfrak{p} \text{ ramified} \\ N\mathfrak{p}^m \leq x}} \log N\mathfrak{p} \\ &\leq \sum_{\substack{\mathfrak{p} \\ \mathfrak{p} \text{ ramified}}} \log N\mathfrak{p} \sum_{N\mathfrak{p}^m \leq x} 1 \\ &\leq 2 \log x \sum_{\substack{\mathfrak{p} \\ \mathfrak{p} \text{ ramified}}} \log N\mathfrak{p} \\ &\leq 2 \log x \log \Delta_L \end{aligned}$$

(We should remark that this estimate would be the same even if C were a union of conjugacy classes.) Also, there are at most n_K distinct pair (\mathfrak{p}, m) such that $N\mathfrak{p}^m = x$, and so

$$\sum_{\substack{\mathfrak{p}, m \\ N\mathfrak{p}^m = x}} \log N\mathfrak{p} \leq n_K \log x.$$

Thus (3.8) yields

$$\psi_C(x) = I_C(x, T) + R_1(x, T), \quad (3.10)$$

where

$$R_1(x, T) \leq 2 \log x \log \Delta_L + n_K \sigma_0 T^{-1} + n_K \log x + R_0(x, T). \quad (3.11)$$

The remainder of this section is devoted to establishing an estimate for $R_0(x, T)$.

So far we allowed σ_0 to be any number > 1 . We now define

$$\sigma_0 = 1 + (\log x)^{-1}. \quad (3.12)$$

While this is only one of many possible choices, it is quite convenient, not least because of the relation $x^{\sigma_0} = ex$.

We now write $R_0(x, T) = S_1 + S_2 + S_3$, where S_1 consists of those terms of (3.9) for which $N\mathfrak{p}^m \leq \frac{3}{4}x$ or $N\mathfrak{p}^m \geq \frac{5}{4}x$, S_2 of those for which $|x - N\mathfrak{p}^m| \leq 1$, and S_3 of the remaining ones. If $N\mathfrak{p}^m \leq \frac{3}{4}x$ or $N\mathfrak{p}^m \geq \frac{5}{4}x$, then

$$\begin{aligned} \left| \log \frac{x}{N\mathfrak{p}^m} \right| &\geq \log \frac{5}{4}, \\ \min \left(1, T^{-1} \left| \log \frac{x}{N\mathfrak{p}^m} \right|^{-1} \right) &\ll T^{-1} \quad \text{for } T \geq 1, \end{aligned}$$

and so

$$S_1 \ll xT^{-1} \sum_{\mathfrak{p}, m} (N\mathfrak{p})^{-m\sigma_0} \log N = xT^{-1} \left[-\frac{\zeta'_K}{\zeta_K}(\sigma_0) \right]. \quad (3.13)$$

To bound this term we use an auxiliary result.

Lemma 3.2. For $\sigma > 1$,

$$-\frac{\zeta'_K}{\zeta_K}(\sigma) \leq -n_K \frac{\zeta'_\mathbb{Q}}{\zeta_\mathbb{Q}}(\sigma).$$

Proof. We have

$$-\frac{\zeta'_K}{\zeta_K}(\sigma) = \sum_{\mathfrak{p}} \frac{\log N\mathfrak{p}}{(N\mathfrak{p})^\sigma - 1}, \quad -\frac{\zeta'_\mathbb{Q}}{\zeta_\mathbb{Q}}(\sigma) = \sum_p \frac{\log p}{p^\sigma - 1}$$

where in the second sum p runs through the rational primes. Now for each prime

ideal \mathfrak{p} , $N\mathfrak{p} = p^k$ for some positive integer k . Thus

$$\frac{\log N\mathfrak{p}}{(N\mathfrak{p})^\sigma - 1} = \frac{k \log p}{p^{k\sigma} - 1} = \frac{k}{p^{(k-1)\sigma} + \dots + 1} \cdot \frac{\log p}{p^\sigma - 1} \leq \frac{\log p}{p^\sigma - 1}.$$

Also, there are at most N_K distinct \mathfrak{p} lying over a given rational prime p , so that

$$-\frac{\zeta'_K}{\zeta_K}(\sigma) \leq n_K \sum_p \frac{\log p}{p^\sigma - 1} = -n_K \frac{\zeta'_\mathbb{Q}}{\zeta_\mathbb{Q}}(\sigma)$$

□

Since

$$-\frac{\zeta'_\mathbb{Q}}{\zeta_\mathbb{Q}}(\sigma) \ll (\sigma - 1)^{-1}$$

for $\sigma > 1$, Lemma 3.2 and (3.13) show that for $T \geq 1$,

$$S_1 \ll n_K x T^{-1} \log x. \quad (3.14)$$

The second sum S_2 consists of those terms \mathfrak{p}^m for which $0 < |N\mathfrak{p}^m - x| \leq 1$.

There are at most $2n_K$ of such \mathfrak{p}^m and since

$$\min\left(1, T^{-1} \left| \log \frac{x}{N\mathfrak{p}^m} \right|^{-1}\right) \leq 1,$$

we obtain

$$S_2 \leq 2n_K \log(x+1) \left(\frac{x}{x-1}\right)^{\sigma_0} \ll n_K \log x. \quad (3.15)$$

The final sum S_3 consists of those terms \mathfrak{p}^m for which $1 < |N\mathfrak{p}^m - x| < \frac{1}{4}x$. Here

we use the estimate

$$\left| \log \frac{x}{n} \right|^{-1} \leq \frac{2n}{|x-n|},$$

valid for $N \geq \frac{1}{2}x$, to obtain

$$\begin{aligned}
S_3 &\ll T^{-1} \log x \sum_{1 < |n-x| < \frac{1}{4}x} \left| \log \frac{x}{n} \right|^{-1} \sum_{\substack{\mathfrak{p}, m \\ N \mathfrak{p}^m = n}} 1 \\
&\ll n_K x T^{-1} \log x \sum_{1 \leq k < \frac{1}{4}x} \frac{1}{k} \\
&\ll n_K x T^{-1} (\log x)^2.
\end{aligned} \tag{3.16}$$

Putting (3.14)-(3.16) together we obtain

$$R_0(x, T) \ll n_K \log x + n_K x T^{-1} (\log x)^2, \tag{3.17}$$

valid for all $x \geq 2$, $T \geq 1$. If we now combine (3.17) with (3.11), we obtain finally the estimate

$$R_1(x, T) \ll \log x \log \Delta_L + n_K \log x + n_K x T^{-1} (\log x)^2, \tag{3.18}$$

valid for all $x \geq 2$, $T \geq 1$, which was the goal of this section. We should mention here that the $\log x \log \Delta_L$ term in (3.18) (which came from the ramified primes) would have been the same even if C were to be the union of any number of conjugacy classes. Let us also note that if $L \neq \mathbb{Q}$, then $n_K \leq n_L \ll \log \Delta_L$, and so the second term on the right side of (3.18) can be absorbed in the first one.

4 Reduction to the case of Hecke L-functions

Our definition (3.5) of $F_C(s)$ was in terms of Artin L-functions corresponding to the (usually nonlinear) characters of $G(L/K)$. In this section we show that $F_C(s)$ can be written in terms of Hecke (abelian) L-functions. This will enable us to obtain much better results on the location and density of the singularities

of $F_C(s)$. The reduction we will use is due to Deuring [Deu35] (later rediscovered by MacCluer [Mac68]). We learned of it from [Mor], and should like to thank J. P. Serre for bringing Moreno's paper to our attention and for supplying the following formulation of Deuring's idea.

In defining $F_C(s)$ by (3.5), we have already selected an element $g \in C$. Let $H = \langle g \rangle$ be the cyclic group generated by g , E the fixed field of H , and let χ denote the irreducible characters of H . Since H is cyclic, the characters χ are one-dimensional. We will retain this notation for the rest of this paper.

Lemma 4.1. We have

$$F_C(s) = -\frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \frac{L'}{L}(s, \chi, L/E). \quad (4.1)$$

Proof. Let $\tau : H \rightarrow \mathbb{C}$ be the class function defined by

$$\tau(h) = \begin{cases} |H| & \text{if } h = g, \\ 0 & \text{if } h \neq g. \end{cases}$$

Then the orthogonality relations for characters of H imply that

$$\tau = \sum_{\chi} \bar{\chi}(g) \chi.$$

Let τ^* denote the class function on G induced by τ , which by direct calculation equals

$$\tau^*(y) = \begin{cases} |C_G(g)| & y \in C, \\ 0 & y \notin C, \end{cases}$$

where $C_G(g)$ is the centralizer of g in G . Now $|C_G(g)||C| = |G|$ so that $\tau^* = f_C$

[see (3.4)]. This implies

$$\sum_{\chi} \bar{\chi}(g)\chi^* = \sum_{\phi} \bar{\phi}(g)\phi,$$

so that for $\operatorname{Re}(s) > 1$ we have

$$F_C(s) = -\frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \frac{L'}{L}(s, \chi^*, L/K). \quad (4.2)$$

But $L(s, \chi^*, L/K) = L(s, \chi, L/E)$, and so (4.1) holds for $\operatorname{Re}(s) > 1$, and therefore (by analytic continuation) for all s . \square

5 Density of zeroes of Hecke L-functions

We have now shown that for $x \geq 2$ and $T \geq 1$, say,

$$\psi_C(x) = I_C(x, T) + R_1(x, T),$$

where $R_1(x, T)$ satisfies (3.18) and

$$I_C(x, T) = -\frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \frac{1}{2\pi i} \int_{\sigma_0 - iT}^{\sigma_0 + iT} \frac{x^s}{s} \frac{L'}{L}(s, \chi, L/E) ds, \quad (5.1)$$

where $\sigma_0 = 1 + (\log x)^{-1}$ and χ runs through the (one-dimensional) irreducible characters of $H = \langle g \rangle$. Our next goal will be to evaluate each of the integrals in (5.1). [This turns out to be more convenient than integrating $F_C(s)$.] To accomplish this we will need some upper bounds on the number of singularities of L'/L .

Since L and E are going to be fixed from now on, we will use $L(s, \chi)$ to denote $L(s, \chi, L/E)$. Also, we let $F(\chi)$ denote the conductor of χ and set

$$A(\chi) = \Delta_E N_{E/Q}(F(\chi)) \quad (5.2)$$

and

$$\delta(\chi) = \begin{cases} 1 & \text{if } \chi = \chi_1, \text{ the principal character,} \\ 0 & \text{otherwise.} \end{cases} \quad (5.3)$$

We recall that for each χ there exist non-negative integers $a = a(\chi)$, $b = b(\chi)$ such that

$$a(\chi) + b(\chi) = n_E, \quad (5.4)$$

and such that if we define

$$\gamma_\chi(s) = \left[\pi^{-\frac{s+1}{2}} \Gamma\left(\frac{s+1}{2}\right) \right]^b \left[\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \right]^a \quad (5.5)$$

and

$$\xi(s, \chi) = [s(s-1)]^{\delta(\chi)} A(\chi)^{s/2} \gamma_\chi(s) L(s, \chi), \quad (5.6)$$

then $\xi(s, \chi)$ satisfies the functional equation

$$\xi(1-s, \bar{\chi}) = W(\chi) \xi(s, \chi), \quad (5.7)$$

where $W(\chi)$ is a certain constant of absolute value 1. Furthermore, $\xi(s, \chi)$ is an entire function of order 1 and does not vanish at $s = 0$, and hence by the Hadamard product theorem we have

$$\xi(s, \chi) = e^{B_1(\chi) + B(\chi)s} \prod_{\rho} \left(1 - \frac{s}{\rho}\right) e^{s/\rho} \quad (5.8)$$

for some constants $B_1(\chi)$ and $B(\chi)$ where ρ runs through all the zeroes of $\xi(s, \chi)$, which are precisely those zeroes $\rho = \beta + i\gamma$ of $L(x, \chi)$ for which $0 < \beta < 1$ [the so-called “nontrivial zeroes” of $L(s, \chi)$]. [We recall that $L(s, \chi)$ and hence $\xi(s, \chi)$ have no zeroes ρ with $\text{Re}(\rho) \geq 1$.] From now on ρ will denote the nontrivial zeroes of $L(s, \chi)$.

Since we are interested in the integrals in (5.1), which involve L'/L , we differentiate (5.6) and (5.8) logarithmically to obtain the important identity

$$\frac{L'}{L}(s, \chi) = B(\chi) + \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right) - \frac{1}{2} \log A(\chi) - \delta(\chi) \left[\frac{1}{s} + \frac{1}{s-1} \right] - \frac{\gamma'_{\chi}}{\gamma_{\chi}}(s), \quad (5.9)$$

valid identically in the complex variable s . A difficulty in the use of this formula is caused by the presence of the constant $B(\chi)$, which depends in an as-yet-undetermined way on χ . However, since $(s-1)^{\delta(\chi)} L(s, \chi)$ is entire, the functional equation (5.7) easily implies the following result which is proved in [Odl77].

Lemma 5.1. With notation as above,

$$\operatorname{Re} B(\chi) = - \sum_{\rho} \operatorname{Re} \frac{1}{\rho}, \quad (5.10)$$

and

$$\begin{aligned} \frac{L'}{L}(s, \chi) + \frac{L'}{L}(s, \bar{\chi}) &= \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{s - \bar{\rho}} \right) - \log A(\chi) \\ &\quad - 2\delta(\chi) \left(\frac{1}{s} + \frac{1}{s-1} \right) - 2 \frac{\gamma'_{\chi}}{\gamma_{\chi}}(s) \end{aligned} \quad (5.11)$$

holds identically in the complex variable s , where ρ runs through the nontrivial zeroes of $L(s, \chi)$.

This lemma will enable us to obtain estimates of both of $B(\chi)$ and of the density of zeroes of $L(s, \chi)$. We should mention, however, that an analog of the above lemma could be proved for general Artin L-functions, but it would contain sums over the possible poles of such L-functions, and these pole terms would prevent us from obtaining an estimate as good as the one below. The purpose of the preceding section's reduction to the case of abelian L-function was to avoid these difficulties.

We first derive some easy auxiliary results.

Lemma 5.2. If $\sigma = \operatorname{Re}(s) > 1$, then

$$\left| \frac{L'}{L}(s, \chi) \right| \ll \frac{n_E}{\sigma - 1}.$$

Proof. A comparison of the Dirichlet series shows that

$$\left| \frac{L'}{L}(s, \chi) \right| \leq -\frac{\zeta'_E}{\zeta_E}(\sigma),$$

and the result follows from Lemma 3.2. □

Lemma 5.3. If $\sigma = \operatorname{Re}(s) > -1/2$ and $|s| \geq 1/8$, then

$$\left| \frac{\gamma'_\chi}{\gamma_\chi}(s) \right| \ll n_E \log(|s| + 2).$$

Proof. This lemma follows from the definition of $\gamma_\chi(s)$ and the fact that

$$\frac{\Gamma'}{\Gamma}(z) \ll \log(|z| + 2)$$

for z satisfying $|z| \geq 1/16$, $\operatorname{Re} z > -1/4$ [WW96, p.251] (cf. Lemma 6.1). □

We now come to the main result of this section. We let $n_\chi(t)$ denote the number of zeroes $\rho = \beta + i\gamma$ of $L(s, \chi)$ with $0 < \beta < 1$, $|\gamma - t| \leq 1$.

Lemma 5.4. For all t we have

$$n_\chi(t) \ll \log A(\chi) + n_E \log(|t| + 2). \quad (5.12)$$

Proof. We evaluate (5.11) at $s = 2 + it$. Lemmas 5.2 and 5.3 imply that

$$\sum_\rho \operatorname{Re} \left(\frac{1}{s - \rho} + \frac{1}{s - \bar{\rho}} \right) \ll \log A(\chi) + n_E \log(|t| + 2). \quad (5.13)$$

But $\operatorname{Re}(s - \rho)^{-1} > 0$ and $\operatorname{Re}(s - \bar{\rho})^{-1} > 0$ since $2 = \operatorname{Re}(s) > \operatorname{Re}(\rho)$ so

$$\begin{aligned} \sum_{\rho} \operatorname{Re} \left(\frac{1}{s - \rho} + \frac{1}{s - \bar{\rho}} \right) &\geq \sum_{\substack{\rho \\ |\gamma - t| \leq 1}} \frac{2 - \beta}{(2 - \beta)^2 + (t - \gamma)^2} \\ &\geq \sum_{\substack{\rho \\ |\gamma - t| \leq 1}} \frac{1}{5} = \frac{1}{5} n_{\chi}(t), \end{aligned}$$

since $1 < 2 - \beta < 2$, which proves the lemma. \square

The bound (5.12) (which is essentially best possible) will be crucial in many of our subsequent arguments. In the case of general Artin L-functions, we could obtain an estimate similar to (5.13), but it would be for the difference of a sum over the zeroes and a similar sum over the poles and the real part of the poles' contribution would be negative.

We now utilize Lemma 5.4 to obtain two additional auxiliary results. We first show that $B(\chi)$ depends mostly on the very small zeroes of $L(s, \chi)$.

Lemma 5.5. For any ε with $0 < \varepsilon \leq 1$ we have

$$B(\chi) + \sum_{\substack{\rho \\ |\rho| < \varepsilon}} \frac{1}{\rho} \ll \varepsilon^{-1} (\log A(\chi) + n_E).$$

Proof. Set $s = 2$ in (5.9) and use lemmas 5.2 and 5.3 to estimate the $L(s, \chi)$ and γ_{χ} terms, respectively. We obtain

$$B(\chi) + \sum_{\rho} \left(\frac{1}{2 - \rho} + \frac{1}{\rho} \right) \ll \log A(\chi) + n_E.$$

Now

$$\left| \frac{1}{2 - \rho} + \frac{1}{\rho} \right| = \frac{2}{|\rho(2 - \rho)|} \leq \frac{2}{|\rho|^2}$$

and so Lemma 5.4 implies

$$\sum_{\substack{\rho \\ |\rho| \geq 1}} \left| \frac{1}{2-\rho} + \frac{1}{\rho} \right| \ll \sum_{j=1}^{\infty} \frac{n_{\chi}(j)}{j^2} \ll \log A(\chi) + n_E.$$

Also, $|2-\rho| \geq 1$, so

$$\sum_{|\rho| < 1} \left| \frac{1}{2-\rho} \right| \ll \log A(\chi) + n_E,$$

and hence

$$B(\chi) + \sum_{\substack{\rho \\ |\rho| < \varepsilon}} \frac{1}{\rho} \ll \sum_{\substack{\rho \\ \varepsilon \leq |\rho| < 1}} \frac{1}{|\rho|} + \log A(\chi) + n_E,$$

which together with Lemma 5.4 completes the proof. \square

Lemma 5.6. If $s = \sigma + it$ with $-1/2 \leq \sigma \leq 3$, $|s| \geq 1/8$, then

$$\left| \frac{L'}{L}(s, \chi) + \frac{\delta(\chi)}{s-1} - \sum_{\substack{\rho \\ |\gamma-t| \leq 1}} \frac{1}{s-\rho} \right| \ll \log A(\chi) + n_E \log(|t| + 2).$$

Proof. We evaluate (5.9) at $\sigma + it$ and $3 + it$ and subtract the resulting relations [in order to eliminate $B(\chi)$] to obtain

$$\begin{aligned} \frac{L'}{L}(s, \chi) - \frac{L'}{L}(3 + it, \chi) &= \sum_{\rho} \left(\frac{1}{s-\rho} - \frac{1}{3+it-\rho} \right) - \frac{\gamma'_{\chi}(s)}{\gamma_{\chi}} \\ &\quad + \frac{\gamma'_{\chi}(3+it)}{\gamma_{\chi}} - \delta(\chi) \left(\frac{1}{s} + \frac{1}{s-1} - \frac{1}{2+it} - \frac{1}{3+it} \right). \end{aligned}$$

We now use Lemmas 5.2 and 5.3 to estimate the $L(3 + it, \chi)$ and the gamma

factors, respectively. We discover that

$$\left| \frac{L'}{L}(s, \chi) + \frac{\delta(\chi)}{s-1} - \sum_{\substack{\rho \\ |\gamma-t| \leq 1}} \frac{1}{s-\rho} \right| \ll n_E \log(|t|+2) + \sum_{\substack{\rho \\ |\gamma-t| > 1}} \left| \frac{1}{s-\rho} - \frac{1}{3+it-\rho} \right| + \sum_{\substack{\rho \\ |\gamma-t| \leq 1}} \left| \frac{1}{3+it-\rho} \right|. \quad (5.14)$$

Since $|3+it-\rho| > 1$ for all ρ and there are $n_\chi(t)$ terms in the last sum, it is $\ll \log A(\chi) + n_E \log(|t|+2)$. For the first sum on the right side of (5.14) we have

$$\begin{aligned} \sum_{\substack{\rho \\ |\gamma-t| > 1}} \left| \frac{1}{s-\rho} - \frac{1}{3+it-\rho} \right| &= \sum_{\substack{\rho \\ |\gamma-t| > 1}} \frac{3-\sigma}{|s-\rho||3+it-\rho|} \\ &\ll \sum_{j=1}^{\infty} \frac{n_\chi(t+j) + n_\chi(t-j)}{j^2} \\ &\ll \log A(\chi) + n_E \log(|t|+2), \end{aligned}$$

and this proves the lemma. \square

6 The contour integral

The next step in the proof is to evaluate $I_C(x, T)$ by evaluating

$$I_\chi(x, T) = \frac{1}{2\pi i} \int_{\sigma_0-iT}^{\sigma_0+iT} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds \quad (6.1)$$

for each character χ of $H = \langle g \rangle$. So far the only condition on T was $T \geq 1$. We now impose the additional requirement that T should not coincide with the ordinate of a zero of any of the $L(s, \chi)$. We also introduce a new parameter, U , which will satisfy $U = j + 1/2$ for some non-negative integer j (eventually we will

let $U \rightarrow \infty$) and define

$$I_\chi(x, T, U) = \frac{1}{2\pi i} \int_{B_{T,U}} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds, \quad (6.2)$$

where $B_{T,U}$ is the positively oriented rectangle with vertices at $\sigma_0 - iT$, $\sigma_0 + iT$, $-U + it$ and $-U - it$. Now $I_\chi(x, T, U)$ can easily be evaluated exactly in terms of the singularities of the integrand as we will show in the next section. In this section, we will show that

$$R_\chi(x, T, U) = I_\chi(x, T, U) - I_\chi(x, T) \quad (6.3)$$

is small.

The remainder $R_\chi(x, T, U)$ may be divided into the vertical integral

$$V_\chi(x, T, U) = \frac{1}{2\pi} \int_{-T}^T \frac{x^{-U+it}}{-U+it} \frac{L'}{L}(-U+it, \chi) dt \quad (6.4)$$

and the two horizontal integrals

$$H_\chi(x, T, U) = \frac{1}{2\pi i} \int_{-U}^{-1/4} \left\{ \frac{x^{\sigma-iT}}{\sigma-iT} \frac{L'}{L}(\sigma-iT, \chi) - \frac{x^{\sigma+iT}}{\sigma+iT} \frac{L'}{L}(\sigma+iT, \chi) \right\} d\sigma, \quad (6.5)$$

$$H_\chi^*(x, T) = \frac{1}{2\pi i} \int_{-1/4}^{\sigma_0} \left\{ \frac{x^{\sigma-iT}}{\sigma-iT} \frac{L'}{L}(\sigma-iT, \chi) - \frac{x^{\sigma+iT}}{\sigma+iT} \frac{L'}{L}(\sigma+iT, \chi) \right\} d\sigma. \quad (6.6)$$

V_χ and H_χ will be estimated by using Lemma 6.2 to bound L'/L . First, however, we prove an auxiliary result about the digamma function.

Lemma 6.1. If $|z+k| \geq 1/8$ for all non-negative integers k , then

$$\frac{\Gamma'}{\Gamma}(z) \ll \log(|z|+2).$$

Proof. If $\operatorname{Re}(z) \geq 1$, this is well known [WW96, p.251]. If $\operatorname{Re}(z) < 1$, then the

recurrence relation

$$\frac{\Gamma'}{\Gamma}(u) = \frac{\Gamma'}{\Gamma}(u+1) - \frac{1}{u}$$

iterated m times shows that

$$\frac{\Gamma'}{\Gamma}(z) = \frac{\Gamma'}{\Gamma}(z+m) - \sum_{k=0}^{m-1} \frac{1}{z+k}$$

for any positive integer m . Choose $m = \lfloor |z| + 2 \rfloor$. Then $\operatorname{Re}(z+m) > 1$, so that

$$\frac{\Gamma'}{\Gamma}(z+m) \ll \log(|z|+2),$$

while $|z+k| \geq 1/8$ for all non-negative integers k implies

$$\sum_{k=0}^{m-1} \frac{1}{z+k} \ll \sum_{k=0}^{m-1} \frac{1}{k+1/8} \ll \log(|z|+2),$$

which proves the lemma. \square

Lemma 6.2. If $s = \sigma + it$ with $\sigma \leq -1/4$, and $|s+m| \geq 1/4$ for all non-negative integers m , then

$$\frac{L'}{L}(s, \chi) \ll \log A(\chi) + n_E \log(|s|+2).$$

Proof. The functional equation (5.7) and the definitions (5.5) and (5.6) imply that

$$\frac{L'}{L}(s, \chi) = -\frac{L'}{L}(1-s, \bar{\chi}) - \log A(\chi) - \frac{\gamma'_\chi}{\gamma_\chi}(1-s) - \frac{\gamma'_\chi}{\gamma_\chi}(s). \quad (6.7)$$

Since $\operatorname{Re}(1-s) \geq 5/4$, we can use Lemma 5.2 to bound $(L'/L)(1-s, \bar{\chi})$. The lemma then follows by an application of Lemma 6.1 to estimate the γ_χ terms. \square

Estimates for $V_\chi(x, T, U)$ and $H_\chi(x, T, U)$ are now very easy to obtain. By the above lemma we have the crude estimates ($U = j+1/2$ so that $|-U+it+m| \geq 1/4$

for all integers m)

$$V_\chi(x, T, U) \ll \frac{x^{-U}}{U} \int_{-T}^T \left| \frac{L'}{L}(-U + it, \chi) \right| dt \ll \frac{x^{-U}}{U} T \{ \log A(\chi) + n_E \log(T + U) \}, \quad (6.8)$$

and

$$\begin{aligned} H_\chi(x, T, U) &\ll \int_{-\infty}^{-1/4} \frac{x^\sigma}{T} (\log A(\chi) + n_E \log(|\sigma| + 2) + n_E \log T) d\sigma \\ &\ll \frac{x^{-1/4}}{T} \{ \log A(\chi) + n_E \log T \} \quad (6.9) \end{aligned}$$

Better estimates can easily be obtained, but would not be too significant, since other error terms will be much larger.

It remains to estimate $H_\chi^*(x, T)$. Lemma 5.6 shows that

$$\frac{L'}{L}(\sigma + iT, \chi) - \sum_{\substack{\rho \\ |\gamma - T| \leq 1}} \frac{1}{\sigma + iT - \rho} \ll \log A(\chi) + n_E \log T$$

if $-1/4 \leq \sigma \leq \sigma_0 = 1 + (\log x)^{-1}$, $x \geq 2$, $T \geq 2$, and a similar estimate holds for L'/L at $\sigma - iT$. Therefore,

$$\begin{aligned} H_\chi^*(x, t) &- \frac{1}{2\pi i} \int_{-1/4}^{\sigma_0} \left\{ \frac{x^{\sigma - iT}}{\sigma - iT} \sum_{\substack{\rho \\ |\gamma + T| \leq 1}} \frac{1}{\sigma - iT - \rho} - \frac{x^{\sigma + iT}}{\sigma + iT} \sum_{\substack{\rho \\ |\gamma - T| \leq 1}} \frac{1}{\sigma + iT - \rho} \right\} d\sigma \\ &\ll \int_{-1/4}^{\sigma_0} \frac{x^\sigma}{T} \{ \log A(\chi) + n_E \log T \} d\sigma \\ &\ll \frac{x}{T \log x} \{ \log A(\chi) + n_E \log T \} \quad (6.10) \end{aligned}$$

To complete our estimate we show that the first integral in (6.10) is not too large.

Lemma 6.3. Let $\rho = \beta + i\gamma$ have $0 < \beta < 1$, $\gamma \neq t$. If $|t| \geq 2$, $x \geq 2$, and

$1 < \sigma_1 \leq 3$, then

$$\int_{-1/4}^{\sigma_1} \frac{x^{\sigma+it}}{(\sigma+it)(\sigma+it-\rho)} d\sigma \ll |t|^{-1} x^{\sigma_1} (\sigma_1 - \beta)^{-1}.$$

Proof. Suppose first that $\gamma > t$. Let B be the rectangle with vertices at $\sigma_1 + i(t-1)$, $\sigma_1 + it$, $-\frac{1}{4} + it$, $-\frac{1}{4} + i(t-1)$, oriented counterclockwise. By Cauchy's theorem,

$$\int_B \frac{x^s}{s(s-\rho)} ds = 0$$

since the integrand has no singularities inside the contour. However, on the three sides of the rectangle other than the segment from $-\frac{1}{4} + it$ to $\sigma_1 + it$, the integrand is majorized by

$$\frac{x^{\sigma_1}}{(|t|-1)(\sigma_1 - \beta)}$$

which proves the result for $\gamma > t$. A similar proof for $\gamma < t$ uses the rectangle with vertices at $\sigma_0 + i(t+1)$, $\sigma_0 + it$, $-\frac{1}{4} + it$, $-\frac{1}{4} + i(t+1)$. \square

The above lemma shows that

$$\begin{aligned} \frac{1}{2\pi i} \int_{-1/4}^{\sigma_0} \frac{x^{\sigma-iT}}{\sigma-iT} \left(\sum_{\substack{\rho \\ |\gamma+T| \leq 1}} \frac{1}{\sigma+iT-\rho} \right) d\sigma &\ll \frac{x^{\sigma_0}}{T} (\sigma_0 - 1)^{-1} n_\chi(-T) \\ &\ll \frac{x \log x}{T} (\log A(\chi) + n_E \log(T)) \quad (6.11) \end{aligned}$$

for $x \geq 2$, $T \geq 2$, and the same estimate holds for the integral involving zeroes ρ with $|\gamma - T| \leq 1$. [Note that if we assume the GRH for $L(s, \chi)$, then we can delete the $\log x$ term in (6.11). Also, even without the GRH we could replace $\log x$ by $\log \log x$ by improving Lemma 6.3.] Therefore we finally obtain

$$H_\chi^*(x, T) \ll \frac{x \log x}{T} (\log A(\chi) + n_E \log T). \quad (6.12)$$

If we now combine (6.8), (6.9), and (6.12), we obtain the main result of this section, namely that

$$\begin{aligned} I_\chi(x, T) - I_\chi(x, T, U) &= -V_\chi(x, T, U) - H_\chi(x, T, U) - H_\chi^*(x, T) \\ &\ll \frac{x \log x}{T} \{\log A(\chi) + n_E \log T\} \\ &\quad + \frac{T x^{-U}}{U} \{\log A(\chi) + n_E \log(T + U)\}. \end{aligned} \quad (6.13)$$

7 The explicit formula

In this section we combine the results of preceding sections in order to obtain an explicit formula for $\psi_C(x)$ in terms of the zeroes ρ .

We first evaluate the integral $I_\chi(x, T, U)$ which was defined by (6.2). We recall that $x \geq 2$, $U = j + 1/2$ for some non-negative integer j , and $T \geq 2$ does not equal the ordinate of any zero of any of the $L(s, \chi)$. By Cauchy's theorem, $I_\chi(x, T, U)$ equals the sum of the residues of the integrand at poles inside $B_{T,U}$. Now if $\chi = \chi_1$, the principal character, then L'/L has a first order pole of residue -1 at $s = 1$, and hence (this term being absent if $\chi \neq \chi_1$) we obtain a contribution of

$$-\delta(\chi)x$$

from the possible pole at $s = 1$. Further, L'/L has a first order pole with residue $+1$ at each nontrivial zero ρ of $L(s, \chi)$ (the ρ 's are counted according to their multiplicity), and so such ρ 's contribute

$$\sum_\rho \frac{x^\rho}{\rho}.$$

In addition, L'/L has first order poles at the so-called trivial zeroes, which are real and nonpositive. In fact, (6.7) shows that L'/L has first order poles at

$s = -(2m - 1)$, $m = 1, 2, \dots$, where the residue is $b(\chi)$, and first order poles at $s = -2m$, $m = 0, 1, 2, \dots$, where the residue is $a(\chi)$. Hence the residues at points s with $\operatorname{Re}(s) < 0$ contribute

$$-b(\chi) \sum_{m=1}^{\lfloor (u+1)/2 \rfloor} \frac{x^{-(2m-1)}}{2m-1} - a(\chi) \sum_{m=1}^{\lfloor U/2 \rfloor} \frac{x^{-2m}}{2m}.$$

The only remaining residue is at $s = 0$, where we have the complication that both x^s/s and L'/L may have first order poles. The Laurent series expansions show that there exist functions $h_1(s)$ and $h_2(s)$ which are analytic at $s = 0$ [$h_2(s)$ depends on χ], such that

$$\frac{x^s}{s} = \frac{1}{s} + \log x + h_1(s)s,$$

and [using (5.9)]

$$\frac{L'}{L}(s, \chi) = \frac{a(\chi) - \delta(\chi)}{s} + r(\chi) + h_2(s)s,$$

where

$$r(\chi) = B(\chi) - \frac{1}{2} \log A(\chi) + \frac{n_E}{2} \log \pi + \delta(\chi) - \frac{b(\chi)}{2} \frac{\Gamma'}{\Gamma} \left(\frac{1}{2} \right) - \frac{a(\chi)}{2} \frac{\Gamma'}{\Gamma}(1). \quad (7.1)$$

Hence the residue at $s = 0$ is

$$r(\chi) + (a(\chi) - \delta(\chi)) \log x.$$

If we now collect all the residue terms, we find that

$$\begin{aligned}
 I_\chi(x, T, U) = & -\delta(\chi)x + \sum_{|\gamma| < T} \frac{x^\rho}{\rho} - b(\chi) + \sum_{m=1}^{\lfloor (u+1)/2 \rfloor} \frac{x^{1-2m}}{2m-1} \\
 & - a(\chi) \sum_{m=1}^{\lfloor u/2 \rfloor} \frac{x^{-2m}}{2m} + r(\chi) + (a(\chi) - \delta(\chi)) \log(x). \quad (7.2)
 \end{aligned}$$

We now let $U \rightarrow \infty$. Then (7.2) and (6.13) give us the explicit formula

$$\begin{aligned}
 I_\chi(x, T) + \delta(\chi)x - \sum_{|\gamma| < T} \frac{x^\rho}{\rho} - r(\chi) - (a(\chi) - \delta(\chi)) \log x \\
 - \frac{n_E}{2} \log(1 - x^{-1}) + \frac{1}{2} (b(\chi) - a(\chi)) \log(1 + x^{-1}) \\
 \ll \frac{x \log x}{T} \{ \log A(\chi) + n_E \log T \}, \quad (7.3)
 \end{aligned}$$

valid for all $x \geq 2$ and all $T \geq 2$ which do not coincide with the ordinate of a zero. If we now let $T \rightarrow \infty$, (7.3) would give us an explicit formula for the inverse Mellin transform

$$\frac{1}{2\pi i} \int_{\sigma_0 - i\infty}^{\sigma_0 + i\infty} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds$$

with no error term. However, for our purposes a cruder version of (7.3) will be more useful.

Theorem 7.1. If $x \geq 2$ and $T \geq 2$, then

$$\begin{aligned}
 \psi_C(x) - \frac{|C|}{|G|} x + S(x, T) \ll \frac{|C|}{|G|} \left\{ \frac{x \log x + T}{T} \log \Delta_L + n_L \log x + \frac{n_L x \log x \log T}{T} \right\} \\
 + \log x \log \Delta_L + n_K x T^{-1} (\log x)^2, \quad (7.4)
 \end{aligned}$$

where

$$S(x, T) = \frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \left\{ \sum_{\substack{\rho \\ |\gamma| < T}} \frac{x^{\rho}}{\rho} - \sum_{\substack{\rho \\ |\rho| < \frac{1}{2}}} \frac{1}{\rho} \right\}. \quad (7.5)$$

[The inner sums in (7.5) are over the nontrivial zeroes ρ of $L(s, \chi)$].

Proof. Lemma 5.5 and (5.4) show that

$$r(\chi) - \sum_{\substack{\rho \\ |\rho| < \frac{1}{2}}} \frac{1}{\rho} \ll \log A(\chi) + n_E,$$

and so

$$I_{\chi}(x, T) + \delta(\chi)x - \sum_{\substack{\rho \\ |\gamma| < T}} \frac{x^{\rho}}{\rho} - \sum_{\substack{\rho \\ |\rho| < \frac{1}{2}}} \frac{1}{\rho} \ll \log A(\chi) + n_E \log x + \frac{x \log x}{T} \{\log A(\chi) + n_E \log T\}.$$

Hence by (5.1) and (6.1) we have for $x \geq 2$, $T \geq 2$ [T not coinciding with the ordinate of any zero ρ of any $L(s, \chi)$]

$$\begin{aligned} I_C(x, T) - \frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \left\{ \delta(\chi)x - \sum_{\substack{\rho \\ |\gamma| < T}} \frac{x^{\rho}}{\rho} - \sum_{\substack{\rho \\ |\rho| < \frac{1}{2}}} \frac{1}{\rho} \right\} \\ \ll \frac{|C|}{|G|} \sum_{\chi} \left\{ \frac{x \log x + T}{T} \log A(\chi) + n_E \log x + \frac{n_E x \log x \log T}{T} \right\} \\ \ll \frac{|C|}{|G|} \left\{ \frac{x \log x + T}{T} \log \Delta_L + n_L \log x + \frac{n_L x \log x \log T}{T} \right\} \end{aligned}$$

since

$$\sum_{\chi} \log A(\chi) = \log \Delta_L$$

by the conductor-discriminant formula, and $n_E \cdot [L: E] = n_L$. Since

$\psi_C(x) = I_C(x, T) + R_1(x, T)$, where $R_1(x, T)$ satisfies (3.18), we obtain the bound of the theorem, provided T does not equal the ordinate γ of some zero $\rho = \beta + i\gamma$.

If, however, $T = \gamma$ for some ρ , then we evaluate (7.4) with T replaced with $T + \varepsilon$ for a very small ε , and let $\varepsilon \rightarrow 0$. The possible discontinuity in the function on the left side comes from zeroes ρ with $T = \gamma$, and since there are $\ll \sum n_\chi(T)$ of them, their contribution can be absorbed in the error term by increasing the constant implied by the \ll notation. \square

The above theorem, which is the main result of this paper, serves to exhibit $\psi_C(x)$ as consisting of the main term $\frac{|C|}{|G|}x$, of $S(x, T)$, and of a relatively small remainder. In the rest of this paper we will be concerned with estimating $S(x, T)$. If we assume the GRH, then a good bound for $S(x, T)$ can be easily given with what we already know. In order to obtain an unconditional result, however, we need to show that the zeroes ρ do not approach close to the line $\operatorname{Re}(s) = 1$.

8 Zero-free regions

In this section we will use the classical method to prove a zero-free region for $\zeta_L(s)$. Since

$$\zeta_L(s) = \prod_{\chi} L(s, \chi) \tag{8.1}$$

and the $L(s, \chi)$ are all analytic for $s \neq 1$, any zero-free region for $\zeta_L(s)$ immediately implies one for each of the $L(s, \chi)$. This approach does have the serious disadvantage that one can often obtain directly with the $L(s, \chi)$ (cf. [DM13, Ch. 14]); in fact, one can essentially replace $\log \Delta_L$ by $\max(\log A(\chi))$ and n_L with n_E in the estimates below. The problem with that result is that in general n_E can be almost as large as n_L and $\max(\log A(\chi))$ almost as large as Δ_L . Finally, we should mention that for a fixed L a better zero-free region can be obtained by more sophisticated methods [Sok68], but the published versions are not explicit as to the dependence on the field L .

Lemma 8.1. There is an absolute, effectively computable constant c_8 such that $\zeta_L(s)$ has no zeroes $\rho = \beta + i\gamma$ in the region

$$\begin{aligned} |\gamma| &\geq (1 + 4 \log \Delta_L)^{-1} \\ \beta &\geq 1 - c_8(\log \Delta_L + n_L \log(|\gamma| + 2))^{-1} \end{aligned}$$

Proof. We have

$$-\frac{\zeta'_L}{\zeta_L} = \sum_{m=1}^{\infty} \alpha(m) m^{-s} \tag{8.2}$$

for $\sigma = \operatorname{Re}(s) > 1$, where $\alpha(m) \geq 0$ for all m . Hence

$$\begin{aligned} \operatorname{Re} \left(-3 \frac{\zeta'_L}{\zeta_L}(\sigma) - 4 \frac{\zeta'_L}{\zeta_L}(\sigma + it) - \frac{\zeta'_L}{\zeta_L}(\sigma + 2it) \right) \\ = \sum_{m=1}^{\infty} \alpha(m) m^{-\sigma} (3 + 4 \cos(t \log m) + \cos(2t \log m)) \geq 0 \end{aligned}$$

by the classical identity

$$3 + 4 \cos \theta + \cos 2\theta = 2(1 + \cos \theta)^2 \geq 0$$

If we now consider the trivial normal extension L of L , then $\zeta_L(s)$ is the Artin L-function associated to the principal character, and if $\gamma_L(s)$ denotes the associated gamma factor then (5.11) shows that

$$2 \frac{\zeta'_L}{\zeta_L}(s) = \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{s - \bar{\rho}} \right) - \log \Delta_L - \frac{2}{s} - \frac{2}{s-1} - 2 \frac{\gamma'_L}{\gamma_L}(s), \tag{8.3}$$

where the summation is over the nontrivial zeroes ρ of $\zeta_L(s)$. We note here that if $\operatorname{Re} s > 1$, then $\operatorname{Re}(s - \rho)^{-1} > 0$ for each zero ρ . If $\rho = \beta + i\gamma$ is some particular

zero with $\gamma \geq (1 + 4 \log \Delta_L)^{-1}$, then we find that for $\sigma > 1$,

$$\begin{aligned} -\frac{\zeta'_L}{\zeta_L}(\sigma) &\leq \frac{1}{\sigma-1} + \frac{1}{\sigma} + \frac{1}{2} \log \Delta_L + \frac{\gamma'_L}{\gamma_L}(\sigma) - \sum_{\rho} \operatorname{Re}(\sigma - \rho)^{-1} \\ &\leq \frac{1}{\sigma-1} + c_9 \log \Delta_L + c_9 n_L, \end{aligned}$$

$$\begin{aligned} -\operatorname{Re} \frac{\zeta'_L}{\zeta_L}(\sigma + 2i\gamma) &\leq \frac{1}{2} \log \Delta_L + \operatorname{Re} \left\{ \frac{1}{\sigma + 2i\gamma - 1} + \frac{1}{\sigma + 2i\gamma} \right\} + \operatorname{Re} \frac{\gamma'_L}{\gamma_L}(\sigma + 2i\gamma) \\ &\leq c_{10} \log \Delta_L + c_{10} n_L \log(|\gamma| + 2), \end{aligned}$$

and

$$-\operatorname{Re} \frac{\zeta'_L}{\zeta_L}(\sigma + i\gamma) \leq c_{11} \log \Delta_L + c_{11} n_L \log(|\gamma| + 2) - \frac{1}{\sigma - \beta},$$

where in the last inequality we have included the contribution of the zero $\rho = \beta + i\gamma$. These inequalities and (8.2) show that for all $\sigma > 1$

$$\frac{4}{\sigma - \beta} < \frac{3}{\sigma - \beta} + c_{12} \{ \log \Delta_L + n_L \log(|\gamma| + 2) \}$$

If we now set $\sigma = 1 + (100c_{12})^{-1} \{ \log \Delta_L + n_L \log(|\gamma| + 2) \}^{-1}$, say, then we obtain the result of the lemma. \square

In addition to Lemma 8.1 we also need information about zeroes $\zeta_L(s)$ very near the real axis. Such information can be obtained by methods very similar to those used above.

Lemma 8.2. If $n_L > 1$ then $\zeta_L(s)$ has at most one zero $\rho = \beta + i\gamma$ in the region

$$\begin{aligned} |\gamma| &\leq (4 \log \Delta_L)^{-1}, \\ \beta &\geq 1 - (4 \log \Delta_L)^{-1} \end{aligned} \tag{8.4}$$

This zero, if it exists, has to be real and simple.

Proof. Identity (8.3) shows that for $1 < \sigma \leq 2$

$$\begin{aligned} \sum_{\rho} \frac{\sigma - \beta}{(\sigma - \beta)^2 + \gamma^2} &= \frac{1}{\sigma - 1} + \frac{1}{2} \log \Delta_L + \frac{\zeta'_L}{\zeta_L}(\sigma) + \frac{1}{\sigma} + \frac{\gamma'_L}{\gamma_L}(\sigma) \\ &\leq \frac{1}{\sigma - 1} + \frac{1}{2} \log \Delta_L \end{aligned} \quad (8.5)$$

since $\zeta'/\zeta \leq 0$ and it is easily verified that

$$\frac{1}{\sigma} + \frac{\gamma'_L}{\gamma_L}(\sigma) = \left(\frac{1}{\sigma} - \frac{n_L}{2} \log \pi \right) + \frac{a(L)}{2} \frac{\Gamma'(\sigma)}{\Gamma(\sigma)} + \frac{b(L)}{2} \frac{\Gamma'(\frac{\sigma+1}{2})}{\Gamma(\frac{\sigma+1}{2})} < 0$$

for $1 < \sigma \leq 1 + (\log 3)^{-1}$. If $\rho = \beta + i\gamma$ is in the region described by (8.4) and $\gamma \neq 0$, then (8.5) gives

$$2 \frac{\sigma - \beta}{(\sigma - \beta)^2 + \gamma^2} \leq \frac{1}{\sigma - 1} + \frac{1}{2} \log \Delta_L,$$

which is false at $\sigma = 1 + (\log \Delta_L)^{-1} \leq 1 + (\log 3)^{-1}$. We similarly obtain a contradiction if there is more than one real zero in our region. \square

If the possible zero described by the above lemma exists, we denote it β_0 and call it the exceptional (Siegel) zero. We also note that if $n_L = 1$ (so that $L = \mathbb{Q}$, $\log \Delta_L = 0$), then ζ_L has no nontrivial zeroes ρ with $|\gamma| < 14$. If β_0 exists, then (8.1) shows that there exists a unique χ_0 such that $L(\beta_0, \chi_0) = 0$. This χ_0 must then be a real character, as $L(\beta_0, \bar{\chi}_0) = \overline{L(\beta_0, \chi_0)} = 0$.

9 Final estimates

We conclude this paper by applying the explicit formula of Theorem 7.1 to estimate $\psi_C(x)$ and $\pi_C(x)$. We start with the GRH estimate for $\psi_C(x)$, which is

the easiest to obtain.

Theorem 9.1. If $\zeta_L(s)$ satisfies the GRH, then

$$\psi_C(x) - \frac{|C|}{|G|}x \ll \frac{|C|}{|G|}x^{\frac{1}{2}} \log x \log \Delta_L x^{n_L} + \log x \log \Delta_L \quad (9.1)$$

for all $x \geq 2$.

Proof. If $\zeta_L(s)$ satisfies the GRH, then so do all of the $L(s, \chi)$. Therefore, for each χ there are no nontrivial zeros ρ with $|\rho| < 1/2$, and so by Lemma 5.4.

$$\begin{aligned} \left| \sum_{|\gamma| < T} \frac{x^\rho}{\rho} + \sum_{|\rho| < \frac{1}{2}} \frac{1}{\rho} \right| &\leq x^{\frac{1}{2}} \sum_{|\gamma| < T} \frac{1}{|\rho|} \\ &\ll x^{\frac{1}{2}} \sum_{j=1}^{\lfloor T \rfloor} \frac{n_X(j)}{j} \\ &\ll x^{\frac{1}{2}} (\log A(\chi) + n_E \log T) \log T, \end{aligned}$$

which together with (7.5) implies

$$S(x, T) \ll \frac{|C|}{|G|}x^{\frac{1}{2}} (\log \Delta_L + n_L \log T) \log T \quad (9.2)$$

for all $T \geq 2$. We choose $T = x^{\frac{1}{2}} + 1$, way, and then (9.2) and (7.4) imply (9.1) for $x \geq 2$. \square

Theorem 9.2. There is an effectively computable positive absolute constant c_{13} such that if

$$x \geq \exp(4n_L(\log \Delta_L)^2) \quad (9.3)$$

then

$$\psi_C(x) = \frac{|C|}{|G|}x - \frac{|C|}{|G|}\chi_0(g) \frac{x^{\beta_0}}{\beta_0} + R(x), \quad (9.4)$$

where

$$|R(x)| \leq x \exp(-c_{13} n_L^{-\frac{1}{2}} (\log x)^{\frac{1}{2}}),$$

and where the second term on the right side of (9.4) occurs only if $\zeta_L(s)$ has an exceptional zero β_0 , and χ_0 is the (real) character of $H = \text{Gal}(L/E) = \langle g \rangle$ for which $L(s, \chi_0, L/E)$ has β_0 as a zero.

Proof. If $\rho = \beta + i\gamma \neq \beta_0$ is a nontrivial zero of one of the $L(s, \chi)$ with $|\gamma| \leq T$, then the unconditional bound of Lemma 8.1 shows that

$$|x^\rho| = x^\beta \leq x \exp\left(-c_{14} \frac{\log x}{\log \Delta_L T^{n_L}}\right)$$

for $x \geq 2$, $T \geq 2$. Further, Lemma 5.4 shows that

$$\sum_{\chi} \sum_{\substack{\rho \\ |\rho| \geq \frac{1}{2} \\ |\gamma| \leq T}} \left| \frac{1}{\rho} \right| \ll \log T \log(\Delta_L T^{n_L}).$$

Also,

$$\sum_{\chi} \sum_{\substack{\rho \neq 1 - \beta_0 \\ |\rho| < \frac{1}{2}}} \left(\left| \frac{x^\rho}{\rho} \right| + \left| \frac{1}{\rho} \right| \right) \ll x^{\frac{1}{2}} \sum_{\chi} \sum_{\substack{\rho \neq 1 - \beta_0 \\ |\rho| < \frac{1}{2}}} \left| \frac{1}{\rho} \right| \ll x^{\frac{1}{2}} (\log \Delta_L)^2,$$

by Lemma 5.4 and the fact that for $\rho \neq 1 - \beta_0$, $|\rho| \geq (4 \log \Delta_L)^{-1}$. (If $\log \Delta_L = 0$, $L = \mathbb{Q}$, and the estimate holds trivially). Finally,

$$\frac{x^{1-\beta_0}}{1-\beta_0} - \frac{1}{1-\beta_0} = x^\sigma \log x \leq x^{\frac{1}{2}} \log x$$

for some σ , $0 \leq \sigma \leq 1 - \beta_0$. Collecting all these estimates gives us

$$S(x, T) - \frac{|C|}{|G|} \chi_0(g) \frac{x^{\beta_0}}{\beta_0} \ll \frac{|C|}{|G|} x \log T \log(\Delta_L T^{n_L}) \exp\left(-\frac{c_{14} \log x}{\log \Delta_L T^{n_L}}\right) + \frac{|C|}{|G|} x^{\frac{1}{2}} (\log \Delta_L)^2. \quad (9.5)$$

We now choose

$$T = \exp(n_L^{-\frac{1}{2}} (\log x)^{\frac{1}{2}} - \log \Delta_L). \quad (9.6)$$

The estimate of the theorem then follows from (9.5) and (7.4). \square

The deduction of Theorem 1.1 and 1.3 from the preceding theorem is now straightforward. We first define the function

$$\theta_C(x) = \sum_{\substack{N_{K/\mathbb{Q}} \mathfrak{p} \leq x \\ \mathfrak{p} \text{ unramified} \\ \left[\frac{L/K}{\mathfrak{p}}\right] = C}} \log(N_{K/\mathbb{Q}} \mathfrak{p}).$$

Since there are at most n_K ideals \mathfrak{p}^m (\mathfrak{p} prime) of a given norm in K ,

$$\sum_{\substack{\mathfrak{p}, m \\ m \geq 2 \\ N_{K/\mathbb{Q}} \mathfrak{p}^m \leq x}} \log(N_{K/\mathbb{Q}} \mathfrak{p}) \ll n_K x^{\frac{1}{2}} \quad (9.7)$$

by an elementary Chebyshev-type estimate. This shows that the estimates of Theorems 9.1 and 9.2 hold when $\psi_C(x)$ is replaced by $\theta_C(x)$. Theorems 1.1 and 1.3 now follow from these estimates for $\theta_C(x)$ by simple partial summation arguments.

We conclude this paper by indicating one way in which the GRH estimate of Corollary 1.2 can be slightly improved. Instead of integrating

$$\frac{1}{2\pi i} \frac{x^s}{s} F_C(x),$$

we can integrate

$$\frac{1}{2\pi i} \left(\frac{y^{s-1} - x^{s-1}}{s-1} \right)^2 F_C(x),$$

where $y > x > 1$, along the contour $B_{T,U}$ of Section 6. We then first let $U \rightarrow \infty$, and then $T \rightarrow \infty$. The integral from $\sigma_0 - i\infty$ to $\sigma_0 + i\infty$ gives us the term we are interested in, i.e.,

$$\sum_{\substack{\mathfrak{p} \\ \left[\frac{L/K}{\mathfrak{p}} \right] = C}} \frac{\log N \mathfrak{p}}{N \mathfrak{p}} r(N \mathfrak{p}; y, x), \quad (9.8)$$

where

$$r(m; y, x) = \begin{cases} \log \frac{m}{x^2} & x^2 \leq m \leq xy, \\ \log \frac{y^2}{m} & xy \leq m \leq y^2, \\ 0 & \text{otherwise,} \end{cases}$$

together with the contribution of the ramified primes and prime powers. By Cauchy's theorem the value of the integral also equals the contribution of the poles of the integrand, which is

$$\frac{|C|}{|G|} \left(\log \frac{y}{x} \right)^2 - \frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \sum_{\rho} \left(\frac{y^{\rho-1} - x^{\rho-1}}{\rho-1} \right)^2, \quad (9.9)$$

where ρ now runs through both the trivial and the nontrivial zeros of $L(s, \chi)$. If we now choose $x = \log \Delta_L$, $y = c_{14}x$, then for c_{14} sufficiently large (and on the assumption of GRH) the main term in (9.9) will dominate both the sum over the zeroes and of the ramified prime and prime power factors, so that (9.8) will have to be nonzero. Hence there will be a prime \mathfrak{p} with $\left[\frac{L/K}{\mathfrak{p}} \right] = C$ and

$$N \mathfrak{p} \leq y^2 \leq c_{14}^2 \log^2 \Delta_L.$$

Bibliography

- [Bom74] Enrico Bombieri. *Le grand crible dans la théorie analytique des nombres*. fr. Astérisque 18. Société mathématique de France, 1974. URL: http://www.numdam.org/item/AST_1987__18__1_0/.
- [Deu35] Max Deuring. “Über den Tschebotareffschen Dichtigkeitssatz”. In: *Mathematische Annalen* 110 (1935), pp. 414–415. DOI: [10.1007/BF01448036](https://doi.org/10.1007/BF01448036).
- [DM13] H. Davenport and H.L. Montgomery. *Multiplicative Number Theory*. Graduate Texts in Mathematics. Springer New York, 2013. ISBN: 9781475759273. URL: <https://books.google.com/books?id=SFztBwAAQBAJ>.
- [Fog61] E Fogels. “On The zeros of Hecke’s L-functions I”. eng. In: *Acta Arithmetica* 7.2 (1961), pp. 87–106. URL: <http://eudml.org/doc/207002>.
- [Gol70] Larry Joel Goldstein. “A Generalization of the Siegel-Walfisz Theorem”. In: *Transactions of the American Mathematical Society* 149.2 (1970), pp. 417–429. ISSN: 00029947. URL: <http://www.jstor.org/stable/1995404> (visited on 10/17/2022).
- [Hei67] H. Heilbronn. “Zeta-functions and L-functions”. In: *Algebraic Number Theory*. Ed. by J.W.S. Cassels, A. Fröhlich, and London Mathematical Society. Acad. Press, 1967, pp. 204–230. ISBN: 9780122689505. URL: https://www.math.arizona.edu/~cais/scans/Cassels-Frohlich-Algebraic_Number_Theory.pdf.
- [Lan71] Serge Lang. “On the zeta function of number fields”. In: *Inventiones Mathematicae* 12.4 (1971), pp. 337–345. DOI: [10.1007/bf01403312](https://doi.org/10.1007/bf01403312).
- [LMO79] J. C. Lagarias, H. L. Montgomery, and A. M. Odlyzko. “A bound for the least prime ideal in the Chebotarev density theorem”. In: *Inventiones Mathematicae* 54.3 (1979), pp. 271–296. DOI: [10.1007/bf01390234](https://doi.org/10.1007/bf01390234).
- [Mac68] Charles MacCluer. “A reduction of the Chebotarev density theorem to the cyclic case”. In: *Acta Arithmetica* 15 (Jan. 1968), pp. 45–47. DOI: [10.4064/aa-15-1-45-47](https://doi.org/10.4064/aa-15-1-45-47).
- [Mor] C.J. Moreno. “An effective Chebotarev density theory”. In: (). (preprint, University of Illinois 1975).
- [Odl77] A.M. Odlyzko. “On Conductors and Discriminants”. In: *Algebraic Number Fields: (L-functions and Galois properties)*. Ed. by A. Fröhlich. Academic Press, 1977, pp. 377–407.
- [Sok68] A.V. Sokolovskii. “Теорема о нулях дзета-функции Дедекинда и расстояние между “соседними” простыми идеалами”. In: *arta arith.* 13 (1968), pp. 321–334. DOI: [10.4064/aa-13-3-321-334](https://doi.org/10.4064/aa-13-3-321-334).
- [Sta74] H.M. Stark. “Some Effective Cases of the Brauer-Siegel Theorem.” In: *Inventiones mathematicae* 23 (1974), pp. 135–152. URL: <http://eudml.org/doc/142257>.
- [Tsc26] N. Tschebotareff. “Die Bestimmung der Dichtigkeit einer Menge von Primzahlen, welche zu einer gegebenen Substitutionsklasse gehören”. In: *Mathematische Annalen* 95 (1926), pp. 191–228. URL: <http://eudml.org/doc/159125>.
- [Wae70] B.L. van der Waerden. *Modern Algebra*. Springer New York, NY, 1970. ISBN: 978-0-387-40624-4. URL: http://www.ru.ac.bd/stat/wp-content/uploads/sites/25/2019/03/104_09_01_van-der-Waerden-B.L.-Modern-Algebra-I-1949.pdf.

- [WW96] E.T. Whittaker and G.N. Watson. *A Course of Modern Analysis*. Cambridge University Press, 1996. ISBN: 9780521588072. URL: http://theory.fi.infn.it/colomo/metodi_old/Whittaker-Watson.pdf.